



## IMMUNE COMPLEXES IN SCHISTOSOMIASIS

### IV — $C_3$ , $C_4$ and $C_{1q}$ characterization and correlation between $C_3$ in serum and circulating IC levels

Ferrucio SANTORO, Daniel BOUT, Daniel CAMUS and André CAPRON

### SUMMARY

$C_3$ ,  $C_4$  and  $C_{1q}$  fractions of complement (C') were investigated in 3% polyethylene glycol (PEG) precipitate of sera from *Schistosoma mansoni* infected patients.  $C_3$  was characterized in 96.8%,  $C_4$  in 90.3% and  $C_{1q}$  in 19.3% of the cases studied. An inversely proportional correlation was demonstrated between  $C_3$  and IC levels in serum.

### INTRODUCTION

The discovery of the  $C_3$  component of the complement (C') in the granular deposits on the glomerular basement membrane of patients with hepatosplenic schistosomiasis<sup>7,15</sup> was very important for the better understanding of the nature of these deposits. IgG and IgM characterizations in glomerulus<sup>7,15</sup> and the recent discovery of a *S. mansoni* specific antigen in a kidney eluate<sup>7</sup> have contributed considerably to the hypothesis that the antigen-antibody complexes are directly responsible for the patients' nephropathy<sup>1,2,4,5,7,11,15</sup>.

In previous studies, the Authors have detected and characterized, by different methods, circulating Immune Complexes (I.C.) in sera from *S. mansoni* infected patients<sup>3,13,14</sup>. In addition to immunoglobulins G, M and E, we have shown the existence of number "4" specific *Schistosoma* antigen in the IC<sup>3</sup>. In a more recent work, the Authors quantified IgG, IgM, IgA and IgE present in IC, obtaining a directly proportional correlation between the quantities of complexed IgM and circulating IC<sup>12</sup>.

In this study, the Authors have investigated the C' in circulating IC, and tried to

correlate the quantities of  $C_3$ ,  $C_4$  and  $C_{1q}$  to the IC levels.

### MATERIAL AND METHODS

#### Sera

Sera from 31 carriers of chronic schistosomiasis were used in this study. Infection by *S. mansoni* was verified by the detection of viable eggs in feces and by serological tests for the diagnosis of schistosomiasis. As control, normal human sera (NHS) of 10 individuals free from any parasitic or microbial infection were used. The anti- $C_{1q}$  serum was obtained from the Hyland Laboratory. Anti- $C_3$  and anti- $C_4$  immunosera were included in the BEHRING plates.

#### Methods

Precipitation with polyethylene glycol (PEG) described by CREIGHTON et al.<sup>6</sup> and modified by the Authors<sup>13</sup>, was used to isolate the IC from the other serum components.

$C_3$  and  $C_4$  were quantified in serum and characterized in 3% PEG precipitated by radial immuno-diffusion as described by MAN-

Service d'Immunologie et de Biologie Parasitaire, Faculté de Médecine et Institut Pasteur, 59000 — LILLE (France) and Laboratorio Central Gonçalo Moniz, FUSEB, SESAB, 40000 — SALVADOR (Bahia) Brasil

Address: Ferrucio SANTORO. Centre d'Immunologie et de Biologie Parasitaire. Institut Pasteur, 20, boulevard Louis XIV — 59012 LILLE CEDEX (France)

CINI et al.<sup>9</sup> on plates supplied by the BEHRING Institute. The same method<sup>9</sup>, was used to quantify the C<sub>1q</sub> present in serum and to characterize it in 3% PEG precipitates.

The C<sub>1q</sub>-<sup>125</sup>I binding test was done according to NYDEGGER et al.<sup>10</sup>.

Results were statistically analyzed by the correlation test establishing a rate of regression over their logarithmical coordinates, and by the chi-square test.

## RESULTS

The presence of C<sub>3</sub>, C<sub>4</sub> and C<sub>1q</sub> in the PEG precipitates of sera from 31 infected patients and 10 controls was studied by radial immunodiffusion<sup>9</sup>. The frequency of these constituents is shown in Table I.

TABLE I

Immune complexes in schistosomiasis. Characterized C<sub>3</sub>, C<sub>4</sub>, and C<sub>1q</sub> in 3% PEG precipitates

Subjects	Number of cases	C3	C4	C1q
Infected patients	31	30 (96.8%)	28 (90.3%)	6 (19.35%)
Controls	10	4 (40%)	1 (10%)	0

A statistically significant difference was observed in the frequency of C<sub>3</sub> and C<sub>4</sub> detected in the 3% precipitates of sera from patients compared to controls ( $p < 0.01$ ).

Regarding C<sub>1q</sub>, although all controls were negative, there is no significant difference, because of its weak positivity in 3% PEG precipitates of serum from infected patients.

C<sub>3</sub>, C<sub>4</sub> and C<sub>1q</sub> in the serum were quantified by radial immunodiffusion<sup>9</sup>, and compared to circulating IC levels evaluated by the C<sub>1</sub>-<sup>125</sup>I binding test<sup>10</sup>. A statistical analysis permitted us to observe an inversely proportional correlation between levels of C<sub>3</sub> and circulating IC ( $r = 0.8238$ ;  $p < 0.001$ ) (Fig. 1).

In our experimental conditions, no correlation was found between C<sub>4</sub> and C<sub>1q</sub>, and circulating IC levels.

## DISCUSSION

The results above show that of 31 infected patients, 30 (97%) presented C<sub>3</sub>, 28 (90%) presented C<sub>4</sub>, and 6 (19%) presented C<sub>1q</sub> in the 3% PEG precipitates. In comparison, the respective percentages in the PEG precipitates of the control group were 40%, 10% and 0%. A significant difference in the frequency of C<sub>3</sub> and C<sub>4</sub> in sera from infected patients compared to controls permits us to conclude that it is probably due to the presence of C<sub>3</sub> and C<sub>4</sub> in circulating IC.

However, our work shows for the first time in schistosomiasis, an inversely proportional correlation between levels of C<sub>3</sub> and circulating IC. In other words, the higher the level of circulating IC, the lower the serum C<sub>3</sub> and vice-versa. A similar relationship was demonstrated by HUNDER & MC DUFFIE<sup>8</sup> and WATTRE et al.<sup>16</sup> in rheumatoid arthritis. This correlation suggests the existence of an important consumption of C<sub>3</sub> by IC in patients' serum. In our experimental conditions, no correlation was observed between C<sub>4</sub> and C<sub>1q</sub>, and IC levels. Two hypotheses may be proposed, excluding an occasional association, for a possible explanation of this phenomenon. Firstly activation of both classical and alternate complement pathways could be admitted. The classical pathway would be assured by the demonstration in this study of complement components in circulating IC from infected patients' serum. The methods of C<sub>1q</sub> binding and complement fixation, used to detect IC, also represent the classical pathway. Alternate pathway activation could also occur. In previous works, the Authors<sup>3,13</sup> have demonstrated the participation of IgE and IgA in circulating IC from schistosome infected patients' serum. When linked to an antigen, these immunoglobulins may activate the alternate pathway. C<sub>1q</sub> and C<sub>4</sub> do not participate in this activation, and are therefore not consumed. In contrast, C<sub>3</sub> is very much implicated.

On the other hand, C<sub>1q</sub> is known to have a very labile linkage with the Fc fragment of a complexed immunoglobulin. In fact, it does not attach itself definitively to Ag-Ab complexes. In this manner the first C<sub>1</sub> component




Figure 1 : IMMUNE COMPLEXES IN SCHISTOSOMIASIS  
CORRELATION BETWEEN  $C_3$  AND IC LEVELS

- ★ confidence limits for the mean
- ★★ confidence limits for the population

may be used several times in the classical complement pathway activation. Likewise  $C_4$ , together with  $C_2$ , participating in a decisive phase of the classical pathway activation, is not fixed definitively to the C' system, when this is linked to the IC. On the contrary,  $C_3$  can be fixed definitively, in any pathway of complement activation.

Although the C' activation can frequently exist in the circulation, as observed in this work, it is possible to foresee a local activation of this system, which would play an important role in renal pathology, specially with the inflammatory phenomena that may be caused by the enzymes freed.

In the future, it could be interesting to carry out a more complete study, including the alternate pathway of activation, which probably has an important role in the mechanisms that take place in the IC deposits.

## RESUMO

**Imuno-Complexos na esquistossomose. IV — Caracterização do  $C_3$ ,  $C_4$  e  $C_{1q}$  e correlação entre as taxas do  $C_3$  Sérico e os IC Circulantes.**

As frações  $C_3$ ,  $C_4$  e  $C_{1q}$  do complemento (C') foram investigadas nos imuno-complexos (IC) circulantes precipitados pelo polietileno-glicol (PEG) do soro de 31 esquistossomóticos.  $C_3$  foi caracterizado em 96,8% dos casos,  $C_4$  em 90,3% e  $C_{1q}$  em 19,3%.

Demonstrou-se uma correlação inversamente proporcional entre as taxas séricas de  $C_3$  e IC.

## ACKNOWLEDGMENTS

We would like to thank Fabienne DERBAUDRENHIEU and Marie-Christine LIEBART for their excellent technical assistance in this work.

SANTORO, F.; BOUT, D.; CAMUS, D. & CAPRON, A. — Immune complexes in schistosomiasis. IV —  $C_3$ ,  $C_4$  and  $C_{1q}$  characterization and correlation between  $C_3$  in serum and circulating IC levels. **Rev. Inst. Med. trop. São Paulo** 19:39-42, 1977.

#### REFERENCES

1. ANDRADE, Z. A. & QUEIROZ, A. C. — Lesões renais na esquistossomose hepatoesplênica. **Rev. Inst. Med. trop. São Paulo** 10: 36-40, 1968.
2. ANDRADE, Z. A.; ANDRADE, S. G. & SADI-GURSKY, M. — Renal changes in patients with hepatosplenic schistosomiasis. **Amer. J. Trop. Med. Hyg.** 20: 77-83, 1971.
3. BOUT, D.; SANTORO, F.; CARLIER, Y. & CAPRON, A. — Imuno-complexos na esquistossomose. III — Caracterização das imunoglobulinas e dos antígenos implicados no IC. **Rev. Inst. Med. trop. São Paulo** 19: 35-38, 1977.
4. BRITO, T. de; BONI, D. de; LOPES, J. D. & SILVA, L. C. da — Kidney biopsy in human schistosomiasis: an ultrastructural study. **Rev. Inst. Med. trop. São Paulo** 11: 62-64, 1969.
5. BRITO, T. de; GUNJI, J.; CAMARGO, M. E.; PENA, D. O. & SILVA, L. C. de — Advanced kidney diseases in patients with hepatosplenic manson's schistosomiasis. **Rev. Inst. Med. trop. São Paulo** 12: 225-235, 1970.
6. CREIGHTON, W. D.; LAMBERT, P. H. & MIESCHER, P. A. — Detection of antibodies and soluble antigen-antibody complexes by precipitation with polyethylene glycol. **J. Immunol.** 111: 1219-1227, 1973.
7. FALCÃO, H. A. & GOULD, D. B. — Immune complex nephropathy in schistosomiasis. **Ann. Inter. Med.** 83: 148-154, 1975.
8. HUNDER, G. G. & McDUFFIE, F. C. — Hypocomplementemia in rheumatoid arthritis. **Amer. J. Med.** 54: 461-472, 1973.
9. MANCINI, G.; CARBONARA, A. O. & HEREMANS, J. F. — Immunochemical quantitation of antigens by single radial immunodiffusion. **Immunochemistry** 2: 235-254, 1965.
10. NYDEGGER, U. E.; LAMBERT, P. H.; GERBER, H. & MIESCHER, P. A. — Circulating immune complexes in the serum in systemic lupus erythematosus and in carriers of hepatitis B antigen. Quantitation by binding to radiolabelled  $C_{1q}$ . **J. Clin. Invest.** 54: 297-309, 1974.
11. QUEIROZ, F. P.; BRITO, E.; MARTINELLI, R. & ROCHA, H. — Nephrotic syndrome in patients with *Schistosoma mansoni* infection. **Amer. J. Trop. Med. Hyg.** 22: 622-628, 1973.
12. SANTORO, F.; BOUT, D.; CAMUS, D.; DESSAINT, J. P. & CAPRON, A. — Immune complexes in schistosomiasis. V — Characterization and quantitation of immunoglobulins present in IC. **Rev. Inst. Med. trop. São Paulo** (In press).
13. SANTORO, F.; BOUT, D. & CAPRON, A. — Imuno-complexos na esquistossomose. II — Dosagem radio-imunológica da ligação do  $C_{1q}$ -I<sup>25</sup> ao IC. **Rev. Inst. Med. trop. São Paulo** 18:293-297, 1976.
14. SANTORO, F.; BOUT, D.; WATTRE, P. & CAPRON, A. — Imuno-complexos na esquistossomose. I — Utilização da fixação do complemento para sua detecção. **Rev. Inst. Med. trop. São Paulo** 18: 152-156, 1976.
15. SILVA, L. C. da; BRITO, T. de; CAMARGO, M. E.; BONI, D. R. de; LOPEZ, J. D. & GUNJI, J. — Kidney biopsy in the hepatosplenic form of infection with *S. mansoni* in man. **Bull. Wld. Hlth. Org.** 42: 907-910, 1970.
16. WATTRE, P.; SANTORO, F.; BOUT, D.; DEFRAZCE, D. & CAPRON, A. — Detection and characterization of circulating immune complexes in rheumatoid arthritis (31 observations). **Biomedecine Express**, 1976 (in press).

Recebido para publicação em 9/2/1976.